arXiv:0711.2117v1 [physics.class-ph] 14 Nov 2007

Fractional Stability

Vasily E. Tarasov

Skobeltsyn Institute of Nuclear Physics,
Moscow State University, Moscow 119991, Russia
E-mail: tarasov@theory.sinp.msu.ru

Abstract

A fractional generalization of variations is used to define a stability of non-integer order. Fractional variational derivatives are suggested to describe the properties of dynamical systems at fractional perturbations. We formulate stability with respect to motion changes at fractional changes of variables. Note that dynamical systems, which are unstable "in sense of Lyapunov", can be stable with respect to fractional variations.

The theory of integrals and derivatives of non-integer order goes back to Leibniz, Liouville, Riemann, Grunwald, and Letnikov [1, 2]. Fractional analysis has found many applications in recent studies in mechanics and physics. The interest in fractional integrals and derivatives has been growing continuously during the last few years because of numerous applications. In a short period of time the list of applications has been expanding. For example, it includes the chaotic dynamics [3, 4], material sciences [5, 6, 7], mechanics of fractal and complex media [8, 9], quantum mechanics [10, 11], physical kinetics [3, 12, 13, 14, 15], long-range dissipation [16, 17], non-Hamiltonian mechanics [18, 19], long-range interaction [22, 23, 24].

In this preprint, we use the fractional generalization of variation derivatives that are suggested in [19]. Fractional integrals and derivatives are used for stability problems (see for example [25, 26, 27, 28, 29]). We consider the properties of dynamical systems with respect to fractional variations [19]. We formulate stability with respect to motion changes at fractional changes of variables (see [20] pages 294-296). Note that dynamical systems, which are unstable "in sense of Lyapunov", can be stable with respect to fractional variations.

Let us consider dynamical system that is defined by the ordinary differential equations. Suppose that the motion of dynamical system is described by the equations

$$\frac{d}{dt}y_k = F_k(y), \quad k = 1, ..., n. \tag{1}$$

Here $y_1, ..., y_n$ be real variables that define the state of dynamical system.

Let us consider the variation δy_k of variables y_k. The unperturbed motion is satisfied to zero value of variation $\delta y_k = 0$. The variation describes that as function $f(y)$ changes at changes of argument y. The first variation describes changes of function with respect to the first power of changes of y:

$$\delta f(y) = D^1_y f(y) dy, \tag{2}$$

where

$$D_y^1 f(y) = \frac{\partial f(y)}{\partial y}.$$

The second variation describes changes of function with respect to the second power of changes of y:

$$\delta^2 f(y) = D_y^2 f(y)(dy)^2. \tag{3}$$

The variation δ^n of integer order n is defined by the derivative of integer order $D_y^n f(y) = \partial^n f/\partial y^n$, such that

$$\delta^n f(y) = D_y^n f(y)(dy)^n.$$

Let us define the variation of fractional order [19] by the equation

$$\delta^\alpha f = D_y^\alpha f \ (\delta y)^\alpha, \tag{4}$$

where D_y^α is a fractional derivative [1, 2] with respect to y.

The fractional variation of order α describes the function $f(y)$ changes with respect to fractional power of variable y changes. The variation of fractional order is defined by the derivative of fractional order.

Let us derive the equations for fractional variations $\delta^\alpha y_k$. We consider the fractional variation of equation (1) in the form:

$$\delta^\alpha \frac{d}{dt} y_k = \delta^\alpha F_k(y), \quad k = 1, ..., n. \tag{5}$$

Using the definition of fractional variation (4), we have

$$\delta^\alpha F_k(y) = [D_{y_l}^\alpha F_k](\delta y_l)^\alpha, \quad k = 1, ..., n. \tag{6}$$

From Eq. (6), and the property of variation

$$\delta^\alpha \frac{d}{dt} y_k = \frac{d}{dt} \delta^\alpha y_k, \tag{7}$$

where $y_k = y_k(t, a)$, we obtain

$$\frac{d}{dt} \delta^\alpha y_k = [D_{y_l}^\alpha F_k] \ (\delta y_l)^\alpha, \quad k = 1, ..., n. \tag{8}$$

Note that in the left hand side of Eq. (8), we have fractional variation of $\delta^\alpha y_k$, and in the right hand side - fractional power of variation $(\delta y_k)^\alpha$.

Let us consider the fractional variation of the variable y_k. Using Eq. (4), we get

$$\delta^\alpha y_k = [D_{y_l}^\alpha y_k] \ (\delta y_l)^\alpha, \quad k = 1, ..., n. \tag{9}$$

For the Riemann-Liouville fractional derivative,

$$D^{\alpha}_{y_l} y_k \neq 0$$

if $k \neq l$. Therefore for simplification of our transformations, we use the fractional derivative as a fractional power of derivative

$$D^{\alpha}_{y_l} y_k = \delta_{kl} D^{\alpha}_{y_l} y_l, \tag{10}$$

where δ_{kl} is a Kronecker symbol. Substituting Eq. (10) into Eq. (9), we can express the fractional power of variations $(\delta y_k)^{\alpha}$ through the fractional variation $\delta^{\alpha} y_k$:

$$(\delta y_k)^{\alpha} = D^{\alpha}_{y_k} y_k \delta^{\alpha} y_k. \tag{11}$$

Substitution of Eq. (11) into Eq. (8) gives

$$\frac{d}{dt}\delta^{\alpha} y_k = \left[D^{\alpha}_{y_l} y_l\right] \left[D^{\alpha}_{y_l} F_k\right] \delta^{\alpha} y_l. \tag{12}$$

Here we mean the sum on the repeated index l from 1 to n. Equation (12) is equations for fractional variations. Let us denote x_k the fractional variations $\delta^{\alpha} y_k$:

$$x_k = \delta^{\alpha} y_k = \left[D^{\alpha}_{y_k} y_k\right] (\delta y_k)^{\alpha}. \tag{13}$$

As a result, we obtain the differential equation for fractional variations

$$\frac{d}{dt} x_k = a_{kl}(\alpha) x_l, \tag{14}$$

where

$$a_{kl}(\alpha) = \left[D^{\alpha}_{y_l} y_l\right] D^{\alpha}_{y_l} F_k. \tag{15}$$

Using the matrix $X^t = (x_1, ..., x_n)$, and $A_{\alpha} = ||a_{kl}(\alpha)||$, we can rewrite Eq. (14) in the matrix form

$$\frac{d}{dt} X = A_{\alpha} X. \tag{16}$$

Equation (16) is a linear differential equation. To define the stability with respect to fractional variations, we consider the characteristic equation

$$Det[A_{\alpha} - \lambda E] = 0 \tag{17}$$

with respect to λ. If the real part $Re[\lambda_k]$ of all eigenvalues λ for the matrix A_{α} are negative, then the unperturbed motion is asymptotically stable with respect to fractional variations. If the real part $Re[\lambda_k]$ of one of the eigenvalues λ of the matrix A_{α} is positive, then the unperturbed motion is unstable with respect to fractional variations.

A system is said to be stable with respect to fractional variations if for every ϵ, there is a δ_0 such that:

$$\|\delta^\alpha y(t_0)\| < \delta_0 \quad => \quad \|\delta^\alpha y(\alpha, t)\| < \epsilon \quad \forall t \in \mathbb{R}_+. \tag{18}$$

The dynamical system is said to be asymptotically stable with respect to fractional variations $\delta^\alpha y(t, \alpha)$ if as

$$t \to \infty, \quad \|\delta^\alpha y(t, \alpha)\| \to 0. \tag{19}$$

The concept of stability with respect to fractional variations (see [20] pages 294-296) is wider than the usual Lyapunov or asymptotic stability. Fractional stability includes concept of "integer" stability as a special case ($\alpha = 1$). A dynamical system, which is unstable with respect to first variation of states, can be stable with respect to fractional variation. Therefore fractional derivatives expand our possibility to research the properties of dynamical systems.

References

[1] S.G. Samko, A.A. Kilbas, O.I. Marichev, *Fractional Integrals and Derivatives Theory and Applications* (Gordon and Breach, New York, 1993)

[2] K.B. Oldham, J. Spanier, *The Fractional Calculus* (Academic Press, New York, 1974)

[3] G.M. Zaslavsky, "Chaos, fractional kinetics, and anomalous transport" Phys. Rep. **371** (2002) 461-580.

[4] G.M. Zaslavsky, *Hamiltonian Chaos and Fractional Dynamics* (Oxford University Press, Oxford, 2005)

[5] *Applications of Fractional Calculus in Physics* (World Scientific, Singapore, 2000)

[6] M. Caputo, *Elasticita e Dissipazione* (Zanichelli, Bologna, 1969) in Italian

[7] R.R. Nigmatullin, "The realization of the generalized transfer equation in a medium with fractal geometry" Phys. Status Solidi B **133** (1986) 425-430; "Fractional integral and its physical interpretation" Theor. Math. Phys. **90** (1992) 242-251.

[8] A. Carpinteri, F. Mainardi, *Fractals and Fractional Calculus in Continuum Mechanics* (Springer, New York, 1997)

[9] V.E. Tarasov, "Continuous medium model for fractal media" Phys. Lett. A **336** (2005) 167-174; "Fractional Fokker-Planck equation for fractal media" Chaos **15** (2005) 023102; "Possible experimental test of continuous medium model for fractal media" Phys. Lett. A **341** (2005) 467-472; "Fractional hydrodynamic equations for fractal media" Ann. Phys. **318** (2005) 286-307; "Wave equation for fractal solid string" Mod. Phys. Lett. B **19** (2005) 721-728; "Dynamics of fractal solid" Int. J. Mod. Phys. B **19** (2005) 4103-4114; "Fractional Chapman-Kolmogorov equation" Mod. Phys. Lett. B **21** (2007) 163-174 (2007).

[10] N. Laskin, "Fractional quantum mechanics" Phys. Rev. E **62** (2000) 3135-3145; "Fractional quantum mechanics and Levy path integrals" Phys. Lett. A **268** (2000) 298-305; "Fractional Schrodinger equation" Phys. Rev. E **66** (2002) 056108.

[11] M. Naber, "Time fractional Schrodinger equation" J. Math. Phys. **45** (2004) 3339-3352.
[12] G.M. Zaslavsky, "Fractional kinetic equation for Hamiltonian chaos" Physica D **76** (1994) 110-122.
[13] A.I. Saichev, G.M. Zaslavsky, "Fractional kinetic equations: solutions and applications" Chaos **7** (1997) 753-764.
[14] G.M. Zaslavsky, M.A. Edelman, "Fractional kinetics: from pseudochaotic dynamics to Maxwell's demon" Physica D **193** (2004) 128-147.
[15] V.E. Tarasov, "Fractional statistical mechanics" Chaos **16** (2006) 033108; "Liouville and Bogoliubov equations with fractional derivatives" Mod. Phys. Lett. B **21** (2007) 237-248.
[16] F. Mainardi, R. Gorenflo, "On Mittag-Leffler-type functions in fractional evolution processes" J. Comput. Appl. Math. **118** (2000) 283-299.
[17] V.E. Tarasov, G.M. Zaslavsky, "Dynamics with low-level fractionality" Physica A **368** (2006) 399-415.
[18] V.E. Tarasov, "Fractional generalization of gradient and Hamiltonian systems" J. Phys. A **38** (2005) 5929-5943; "Fractional generalization of gradient systems" Lett. Math. Phys. **73** (2005) 49-58.
[19] V.E. Tarasov, "Fractional variations for dynamical systems: Hamilton and Lagrange approaches" J. Phys. A **39** (2006) 8409-8425.
[20] V.E. Tarasov, "Fractional derivative as fractional power of derivative" International Journal of Mathematics. Vol.18. No.3. (2007) pp.281-299.
[21] V.E. Tarasov, "Psi-series solution of fractional Ginzburg-Landau equation" J. Phys. A **39** (2006) 8395-8407.
[22] N. Laskin, G.M. Zaslavsky, "Nonlinear fractional dynamics of lattice with long-range interaction" Physica A **368** (2005) 38-54.
[23] V.E. Tarasov, G.M. Zaslavsky, "Fractional dynamics of coupled oscillators with long-range interaction" Chaos **16** (2006) 023110; "Fractional dynamics of systems with long-range interaction" Commun. Nonlin. Sci. Numer. Simul. **11** (2006) 885-898.
[24] V.E. Tarasov "Map of discrete system into continuous" J. Math. Phys. **47** (2006) 092901. "Continuous limit of discrete systems with long-range interaction" J. Phys. A **39** (2006) 14895-14910.
[25] S. Momani, S.B. Hadid "Lyapunov stability solutions of fractional integrodifferential equations" Int. J. Math. Math. Sci. **47** (2004) 2503-2507.
[26] S.B. Hadid, J.G. Alshamani, "Liapunov stability of differential equations of noninteger order" Arab J. Math **7** (1986) 5-17.
[27] Y.Q. Chen, K.L. Moore, "Analytical stability bound for a class of delayed fractional-order dynamic systems" Nonlinear Dynamics **29** (2002) 191-200.
[28] T.D. Khusainov, "Stability analysis of a linear-fractional delay system" Differ. Eq. **37** (2001) 1184-1188.
[29] D. Matignon, "Stability result on fractional differential equations with applications to control processing" In: IMACS - SMC Proceeding, Lille, France, 1996, pp. 963-968.